




Supplemental Digital Content 1. Incidence of early nausea: Cumulative incidence 0-6h postoperatively




Supplemental Digital Content 2. Incidence of early vomiting: Cumulative incidence 0-6h postoperatively




Supplemental Digital Content 3. Incidence of early PONV (any nausea and/or vomiting): Cumulative incidence 0-6h postoperatively




Supplemental Digital Content 4. Incidence of late nausea: Cumulative incidence 0-24h postoperatively




Supplemental Digital Content 5. Incidence of late vomiting: Cumulative incidence 0-24h postoperatively




Supplemental Digital Content 6. Incidence of late PONV (any nausea and/or vomiting): Cumulative incidence 0-24h postoperatively




Supplemental Digital Content 7. Incidence of early nausea: Cumulative incidence 0-6h postoperatively. Inhalational anesthetics, without anti-emetic prophylaxis




Supplemental Digital Content 8. Incidence of early vomiting: Cumulative incidence 0-6h postoperatively. Inhalational anesthetics, without anti-emetic prophylaxis




Supplemental Digital Content 9. Incidence of early PONV (any nausea and/or vomiting): Cumulative incidence 0-6h postoperatively. Inhalational anesthetics, without anti-emetic prophylaxis




Supplemental Digital Content 10. Incidence of late nausea: Cumulative incidence 0-24h postoperatively. Inhalational anesthetics, without anti-emetic prophylaxis



Supplemental Digital Content 11. Incidence of late vomiting: Cumulative incidence 0-24h postoperatively. Inhalational anesthetics, without anti-emetic prophylaxis



Supplemental Digital Content 12. Incidence of late PONV (any nausea and/or vomiting): Cumulative incidence 0-24h postoperatively. Inhalational anesthetics, without anti-emetic prophylaxis



Supplemental Digital Content. List of references

6. Goll V, Akca O, Greif R, Freitag H, Arkilic CF, Scheck T, Zoeggeler A, Kurz A, Krieger G, Lenhardt R, Sessler DI: Ondansetron is no more effective than supplemental intraoperative oxygen for prevention of postoperative nausea and vomiting. *Anesth Analg* 2001; 92: 112-7
 7. Greif R, Laciny S, Rapf B, Hickle RS, Sessler DI: Supplemental oxygen reduces the incidence of postoperative nausea and vomiting. *Anesthesiology* 1999; 91: 1246-52
 27. McKeen DM, Arellano R, O'Connell C: Supplemental oxygen does not prevent postoperative nausea and vomiting after gynecological laparoscopy. *Can J Anaesth* 2009; 56: 651-7
 29. Simurina T, Mraovic B, Mikulandra S, Sonicki Z, Sulen N, Dukic B, Gan TJ: Effects of high intraoperative inspired oxygen on postoperative nausea and vomiting in gynecologic laparoscopic surgery. *J Clin Anesth* 2010; 22: 492-8
 30. Thibon P, Borgey F, Boutreux S, Hanouz JL, Le Coutour X, Parienti JJ: Effect of perioperative oxygen supplementation on 30-day surgical site infection rate in abdominal, gynecologic, and breast surgery: the ISO2 randomized controlled trial. *Anesthesiology* 2012; 117: 504-11
 57. Bhatnagar S, Mishra S, Gupta M, Singhal AK, Mandal AS: Effects of different concentrations of intraoperative supplemental oxygen on post-operative nausea and vomiting (PONV) in patients under going modified radical mastectomy The Internet Journal of Anesthesiology 2005; 9:12
 59. Joris JL, Poth NJ, Djamadar AM, Sessler DI, Hamoir EE, Defechereux TR, Meurisse MR, Lamy ML: Supplemental oxygen does not reduce postoperative nausea and vomiting after thyroidectomy. *Br J Anaesth* 2003; 91: 857-61
 65. Piper SN, Rohm KD, Boldt J, Faust KL, Maleck WH, Kranke P, Suttner SW: Inspired oxygen fraction of 0.8 compared with 0.4 does not further reduce postoperative nausea and vomiting in dolasetron-treated patients undergoing laparoscopic cholecystectomy. *Br J Anaesth* 2006; 97: 647-53
 67. Purhonen S, Niskanen M, Wustefeld M, Hirvonen E, Hynynen M: Supplemental 80% oxygen does not attenuate post-operative nausea and vomiting after breast surgery. *Acta Anaesthesiol Scand* 2006; 50: 26-31
 68. Purhonen S, Turunen M, Ruohoaho UM, Niskanen M, Hynynen M: Supplemental oxygen does not reduce the incidence of postoperative nausea and vomiting after ambulatory gynecologic laparoscopy. *Anesth Analg* 2003; 96: 91-6
 71. Turan A, Apfel CC, Kumpch M, Danzeisen O, Eberhart LH, Forst H, Heringhaus C, Isselhorst C, Trenkler S, Trick M, Vedder I, Kerger H: Does the efficacy of supplemental oxygen for the prevention of postoperative nausea and vomiting depend on the measured outcome, observational period or site of surgery? *Anaesthesia* 2006; 61: 628-33
-