

# Regimens

## Acetaminophen

| multiple dose |   |  |  |
|---------------|---|--|--|
| paracetamol   | 1g /6h po<br>1g /4h po<br>1g /6h iv<br>1.3g /8h pr<br>1g /6h pr |  |  |
| propacetamol  | 2 g /6h iv  |  |  |

## Non-steroidal anti-inflammatory drugs

|               | multiple dose  | continuous infusion            | single dose             |
|---------------|--|--------------------------------|-------------------------|
| diclofenac    | 50-100 mg /8h iv<br>50-100 mg /8h pr<br>75 mg /12h pr<br>75 mg /12h im | 2 mg/kg /h iv                  | 100 mg pr<br>75 mg iv   |
| ibuprofen     | 500 mg /8h pr  |                                | 1600 mg po              |
| indomethacin  |  |                                | 75 mg po                |
| ketoprofen    | 100 mg /8h iv<br>100 mg /12h iv<br>50 mg /6h iv<br>100 mg /12h im      |                                |                         |
| dexketoprofen | 50 mg /12h im  |                                |                         |
| ketorolac     | 15-30 mg /6h iv<br>15 mg /3h iv<br>15-30 mg /6h im<br>10 mg /4h im     | 2-5 mg /h iv<br>1.5-3 mg /h im | 30-60 mg iv             |
| meloxicam     |  |                                | 15 mg pr                |
| naproxen      |  |                                | 550 mg po               |
| piroxicam     |  |                                | 40 mg iv                |
| tenoxicam     |  |                                | 20-40 mg iv<br>40 mg im |

## Selective cyclo-oxygenase 2 inhibitors

|            | multiple low dose | multiple high dose             | single dose |
|------------|-------------------|--------------------------------|-------------|
| celecoxib  |                   |                                | 200 mg po   |
| parecoxib  | 20 mg /12h iv     | 40 mg /12h iv<br>40 mg / 6h iv |             |
| rofecoxib  | 25 mg /12h po     | 50 mg /12h po                  |             |
| valdecoxib | 20 mg /12h po     | 40 mg /12h po                  |             |

iv: intravenously. im: intramuscularly. pr: rectally. po: orally.