Fig. D: Time to onset of motor block

Review: Clonidine as adjuvant for peripheral nerve and plexus blockades
Comparison: 03 Block times
Outcome: 03 Onset of motor block

Study or sub-category	Treatment N	Treatment Mean (SD)	Control N	Control Mean (SD)	Weight %	WMD (random) 95% CI
Adnan	13	7.10(3.70)	15	6.40(4.90)	20.55	0.70 [-2.49, 3.89]
Duma-Bupivacaine	20	23.30(16.49)	18	18.90(17.62)	5.24	4.40 [-6.49, 15.29]
Duma-Levobupivacaine	20	27.60(46.29)	20	25.30(29.78)	1.25	2.30 [-21.82, 26.42]
El Saied	23	17.90(1.70)	23	16.30(1.40)	27.51	1.60 [0.70, 2.50]
Fang-Bupivacaine	15	22.50(4.00)	15	23.80(4.50)	21.06	-1.30 [-4.35, 1.75]
Fang-Lidocaine	15	11.10(2.60)	15	15.00(3.20)	24.39	-3.90 [-5.99, -1.81]
Fang-Ropivacaine	15	15.00(0.00)	15	21.30(4.30)		Not estimable
Total (95% CI)	121		121		100.00	-0.38 [-3.14, 2.38]

Test for heterogeneity: Chi² = 24.59, df = 5 (P = 0.0002), I² = 79.7%
Test for overall effect: Z = 0.27 (P = 0.79)