

Federal Office of Public Health FOPH  
Public Health Directorate Communicable  
Diseases Division

Schwarzenburgstrasse 157  
3003 Bern  
Switzerland

Geneva, December 23, 2022

## Swiss national SARS-CoV-2 genomic and variants surveillance program: report of the month of November 2022

Geneva Centre for  
Emerging Viral Diseases

Division of Infectious  
Diseases

Department of Medicine

Laboratory of virology

Division of Laboratory  
Medicine

Diagnostic Department

### 1. Summary

In November COVID-19 cases numbers apparently decreased in Switzerland, with vast majority of cases still caused by BA.5 or its sublineages.

Approximately 2.6% of the total number of cases identified in Switzerland in November were sequenced by the Surveillance program, yielding over 2'100 sequences. The majority of the sequences in Switzerland belong to the BQ.1 sublineage of BA.5.

The currently circulating BA.5 variants are at least partially resistant to all the monoclonal antibody therapies used in Switzerland. In particular they are resistant to Evusheld®, which is composed of tixagevimab (complete resistance) and cilgavimab (partial resistance); and sotrovimab, for which the clinical effectiveness is unclear.

BA.4 and 5 sublineages with Spike 346 mutations enabling complete escape from cilgavimab (thus, complete escape from Evusheld®) are growing in frequency. In November, these resistant variants represented over 79% of the sequences identified in Switzerland. Preliminary sequence data for December shows a proportion of nearly 90%. **All available monoclonal antibodies available in Switzerland are expected to be ineffective against most circulating SARS-CoV-2 viruses as of the time of this report.**

The escape from immunity derived from vaccination or previous exposure by these subvariants is weaker than the escape seen against the monoclonal antibodies, but still substantial and it continues to grow. While the bivalent BA.1 booster is not ideal and would benefit from a further update, available data suggests that it will perform significantly better than the previous vaccines against the currently circulating variants. Most importantly these subvariants have so far not resulted in a large increase of hospitalizations.

## **2. Description of the Swiss national SARS-CoV-2 genomic and variants surveillance program.**

The overall goal of the program is to provide epidemiological trends and to highlight meaningful observations.

Because greater transmissibility and/or immune escape potential of the different VOCs and VOIs can result in new surges in COVID-19 numbers despite the vaccination campaign, this program aims to closely monitor each variant displaying mutations known to be linked with either increased transmissibility or immune escape potential.

Currently, 12 diagnostic laboratories are participating in the program, including university hospital centres in Switzerland (Geneva, Lausanne, Bern, Basel, Zurich, Ticino), in addition to private laboratories (Viollier; Medisupport including Dianalabs, Polyanalytic, Dianalabs Valais, Proxilab and Bioanalytica; Labor Team W, Risch), cantonal-based laboratories (Hôpital du Valais – Institut Central), Spital Region Oberrhein (Bern, Solothurn, Aargau, Luzern), and 2 high-throughput sequencing platforms (Health 2030 Genome Centre in Geneva, Genomics Facility Basel run by ETH Zürich and University of Basel). In the month of October, sequencing in Geneva was partially funded by the EU grant for the COVICIS project (<https://covicis.eu/>).

Processed sequencing data are shared openly within 14 days from positive PCR result through the GISAID platform (<https://www.gisaid.org>) and eventually through the Swiss Pathogen Surveillance Platform (SPSP). The centralized analysis of this National Surveillance will be performed by the groups of Pr. Neher, Pr. Stadler and Dr. Althaus, where variants of concern are counted, analyzed and all sequences scanned for new variants with potential changes in antibody-Spike interactions (<https://nextstrain.org/groups/swiss>, <https://covariants.org/per-country>, <https://cov-spectrum.ethz.ch>). This work is done in close collaboration with the Swiss National COVID-19 Science Task Force and the Swiss Institute of Bioinformatics (SIB).

In order to complement the genomic surveillance based on patient samples, sequencing of SARS-CoV-2 in wastewater samples is also performed. Samples are collected daily in six wastewater treatment plants (WWTP), under the coordination of Eawag. Up to 50 samples per week over the first 26 weeks have been performed. The sequencing and analysis of these samples, including detection of variants, is done under the coordination of Prof Niko Beerenwinkel. It started in December 2020 for Lausanne and Zurich, and in February 2021 for all six WWTP (<https://bsse.ethz.ch/cbg/research/computational-virology/sarscov2-variants-wastewater-surveillance.html>).

Immunological characterization of the variants within the surveillance program is coordinated by Professor Trono's team at EPFL.

This report has been produced by Erik Boehm, Pauline Vetter, Marc Friedli, Samuel Cordey, Richard Neher, Christian Althaus, Martina Reichmuth, Cornelius Römer, Niko Beerenwinkel, David Dreifuss, Chaoran Chen, Tanja Stadler, Priscilla Turelli, Didier Trono, Emma Hodcroft, Erik Studer, Oluwaseun Oyewole, and Laurent Kaiser. The list of the participants and collaborators of the program can be found at the end of this report in the appendix.

This report covers the period of 31 October to 27 November, 2022 (weeks 44, 45, 46, 47). All data presented in this report are based on the sampling date.

### **3. Variants of concern (VOCs), variant of interest (VOI) and other surveilled variants: brief summary and special focus**

Five variants and their sub-lineages are considered VOCs by the WHO, B.1.1.7 (Alpha), B.1.351 (Beta), P.1 (Gamma), B.1.617.2 (Delta), and B.1.1.529 (Omicron). Worldwide, all VOCs except Omicron have essentially disappeared from samples collected within the last 30 days (<https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19---24-august-2022>).

#### **Omicron**

The Omicron VOC (B.1.1.529) is characterized by a high divergence in the spike protein, which has allowed it to substantially escape immunity conferred by vaccination (using the original Wu-1 sequence) and prior infection with pre-Omicron variants. This VOC currently has 3 sublineages that still have significant circulation: BA.2, 4, and 5, all of which have further “sub-sublineages”. Despite all being considered “Omicron”, these sublineages may differ from each other (in terms of mutation counts) more than the earlier VOCs differed from the original Wu-1 strain.

Notably, this is the first VOC to have subvariants causing multiple successive waves, arguing for treating Omicron subvariants as distinct VOCs. These sublineages have successively replaced each other, with BA.5 currently being dominant.

While BA.5 replaced BA.2, highly derived BA.2 sublineages have appeared, in particular, BA.2.75. While the growth of this variant has largely been confined to a few countries, some of its subvariants (see below) show signs of more growth potential.

Subvariants of BA.4 and BA.5 containing additional mutations, particularly at position 346 or 444 of the spike protein (such as BQ.1 and BA.4.6), seem to confer a growth advantage and enable complete escape from cilgavimab, rendering it fully resistant to Evusheld®. In particular, subvariants with the R346T mutation have strong signs of growth and are dominant or nearly dominant in many western countries. Notably BQ.1 and sublineages thereof (which contain spike 346 or 444 mutations) seem to be the currently dominant variants in Switzerland.

BA.2.75.2, which has a spike 486 mutation enabling complete escape from tixagevimab like BA.5, is also spawning subvariants. BA.2.75 subvariants also seem to be picking up spike 346 or 444 mutations enabling complete escape from cilgavimab.

A recombination event between a BA.2.10 sublineage (BJ.1) and a BA.2.75 sublineage (BM.1.1.1) has produced the XBB lineage, which is also showing strong signs of growth and immune escape. Notably, recombinants between BA.5 and BA.2.75 sublineages have also been noted (XBF and XBD), and have strong immune escape properties as well.

#### **Detection**

All sub-lineages are still detected by RT-PCR tests, and all except BA.2 exhibit S-gene target failure with the Roche PCR assays regularly used in Switzerland. Given the current virus circulating, the absence of S-gene target failure is currently a good proxy for BA.2.75 infection. Likewise, its presence is indicative of a likely BA.5 (or BA.5 subvariant) infection. Further discrimination between subvariants is not feasible at this time by any method other than genomic sequencing.

Antigenic tests are still able to detect these variants, and sensitivity to the currently circulating variants is relatively unchanged relative to the initial virus. There is some evidence that sensitivity may decrease depending on the patient’s immune status, which may confound results. There is no evidence that the new subvariants pose any particular detection challenges to these tests.

#### **Immune escape**

Extensive data demonstrates that Omicron variants are substantially able to evade neutralizing antibodies (nAbs) from non-Omicron infections and after 2-3 doses of vaccine. Escape from monoclonal antibodies is extensive and is covered by the “Therapeutic intervention effectiveness” section.

Data suggests low neutralization of BA.2.75 and BA.4/5 by sera from BA.1 and BA.2 vaccine-breakthrough infections. For BA.5, this was shown to lead to decreased protection against hospitalization relative to BA.2 infections, although this difference was reduced after 3 vaccine doses, and it was nearly absent after 4 doses. BA.4 and BA.5 sublineages containing further escape mutations, in particular BQ.1\* and XBB, have been shown to be substantially less neutralized than earlier variants by patient sera, even if the sera comes from people exposed to “basic” BA.4 or BA.5 as well as people who have been boosted by Bivalent vaccines. Similarly, the BA.2.75.2 sublineage neutralization titers were lower than the neutralization titers for BA.5. Furthermore, these sublineages completely escape neutralization by Evusheld® and have substantially reduced neutralization by sotrovimab. Unlike the monoclonal antibody situation, neutralization capacity of patient sera against new variants is expected to gradually decline rather than suffer sharp drops in or complete loss of efficacy. Still, it is clear that additional updates would be very beneficial.

While all Omicron sublineages largely escape humoral immunity from pre-omicron vaccines and infections, cell mediated immunity remains mostly intact. Efficacy preventing hospitalization and death is reduced but remains high after three doses and is even higher after four doses. The efficacy of four doses at preventing symptomatic disease is relatively high (>60%), but remains poor (~30%) at preventing any infection.

### **Severity**

A multitude of clinical, *in vitro* and *in vivo* studies indicate that Omicron BA.1 and BA.2 cause intrinsically milder disease. Importantly, BA.1,2, and 3 use a TMPRSS2-independent entry pathway, and exhibit less cleavage of the spike protein and corresponding cell to cell fusion. This suggests an altered tropism that may favor replicating in the upper rather than lower respiratory tract, and is a partial explanation for observations of the infections being less severe. BA.4 and BA.5 have apparently regained the ability to use TMPRSS2 mediated entry pathways and exhibit heightened cell to cell fusion relative to BA.1&2 (although it is still not quite as high as that of earlier variants), suggesting that their intrinsic severity may be closer to that of previous variants.

Other studies of chimeric viruses (the ancestral virus with its spike replaced by an Omicron Spike) in mice suggest that the reduced severity is not due to properties of the spike protein and receptor usage/entry pathway. Notably, all Omicron lineages contain a T9I mutation in the E gene, which has been implicated in reduced severity.

There are conflicting animal studies with regards to the question of intrinsic severity, and severity determinations are complicated by an over representation of reinfections/vaccine breakthroughs, which are expected to be mild due to the protective effect of prior vaccination/exposure.

There is currently no evidence that the severity of the new subvariants (such as XBB, BQ.1.1, and BA.2.75.2) has significantly changed.

### **Therapeutic intervention effectiveness**

All sublineages display complete escape from combination of casirivimab/imdevimab. A matched cohort study found a noticable clinical benefit of sotrovimab treatment during a BA.1 wave. Both *in vitro* and *in vivo* data suggests that sotrovimab is even less effective against BA.2, 4 and 5. Indeed a study in Qatar failed to find any beneficial effect of sotrovimab treatment in the context of BA.2 infections. In contrast a study done in the USA found a beneficial effect for both BA.1 and BA.2, although it had significant limitations specifically regarding its BA.2 conclusions. Currently, there is conflicting data regarding the efficacy of sotrovimab against BA.2.75, with reports variously indicating that its sensitivity is relatively unchanged or decreased relative to BA.5. Multiple studies have reported that BQ.1.1 has a very strong or complete escape against Sotrovimab, even compared to BA.2 and BA.5

Similarly, *in vitro* data suggests that both antibody components of Evusheld® (tixagevimab and cilgavimab) will have significantly reduced neutralization against BA.4/5, but one component (cilgavimab) should retain efficacy against BA.2. Additional spike 346 mutations seen in BA.4/5 sublineages lead to complete escape from cilgavimab, and it is likely that cilgavimab has lost any efficacy that it still had as these sublineages have proliferated. Notably, these lineages include BQ.1.1 which is also reported to be highly resistant to Sotrovimab, and is outcompeting its parent lineages.

As sotrovimab and tixagevimab/cilgavimab are still being used, additional mutations causing escape from them will thus be closely followed (Table 1, and section 7).

In addition to mAbs, there are other antiviral treatments, such as the 3CL-like protease inhibitor Paxlovid® (PF-07321332, nirmatrelvir/ritonavir) or RNA nucleotide analogues (such as molnupiravir). Patients taking Paxlovid have been noted to have a higher “rebound” frequency than those who do not take the drug. Importantly, while viral loads may go up when the drug is discontinued, the drug still efficiently limits viral replication when given and is very effective at avoiding hospitalization and death. Resistance mutations to Paxlovid® have been noted. Given the low proportion of the population that receives Paxlovid®, it is there appears to be little selection pressure for these. Indeed, known escape mutations against Paxlovid® are currently very rare.

Resistance to molnupiravir comes at the cost of significantly reduced viral fitness, so limited use is unlikely to lead to resistant strains. Preliminary data confirms that molnupiravir, Paxlovid®, and remdesivir all retain full *in vitro* efficacy against Omicron sub-lineages.

AA position	World	Europe	Switzerland
<b>Sotrovimab</b>			
337	0.04	0.03	0
340	0.06	0.07	0.09
356	4.41	4.73	4.95
371	95.23	93.85	99.29
377	0.03	0.02	0
449	0.01	0	0
476	0.09	0.06	0
494	0.25	0.25	0.66
<b>Cilgavimab</b>			
346	50.14	56.39	66.4
444	36.19	39.05	47.36
445	5.34	3.45	4.43
446	11.7	9.87	10.84
447	0	0	0
448	0.01	0	0
450	2.6	3.03	1.93
452	84.9	86.51	88.12
484	96.41	95.34	99.72
494	0.25	0.25	0.66
655	98.78	98.66	99.67
<b>Paxlovid®</b>			
54	0.02% (48)	0	0
138	0.00% (2)	0	0
144	0.00% (2)	0	0
165	0.00% (2)	0	0
166	0.00% (5)	0	0
167	0.00% (2)	0	0
172	0.00% (3)	0	0
192	0.00% (8)	0	0
194	0.01% (17)	0.00% (4)	0
305	0.00% (2)	0	0

*Table 1: Frequency of mutations at residues linked (by deep mutations scanning or other experimental results) to escape from sotrovimab, cilgavimab, or Paxlovid®, November 2022. Note the high proportion of mutations at multiple escape sites for cilgavimab, compared to the low number of mutations at sites leading to escape from Paxlovid.*

The circulation of specific mutations leading to decreased effectiveness of known therapeutics available for clinical use in Switzerland will be surveilled (see Section 7), but their prevalence remains low.

Notably, there has been a substantial and worrying increase in Europe and the rest of the world in mutations leading to escape from Cilgavimab (particularly at position 346 and 444). In November, variants with resistance mutations expected to lead to complete escape from both Cilgavimab and Tixagevimab represented over 79% of the sequences identified in Switzerland. Preliminary data for December (with data available as of 22-12) shows a proportion of nearly 90%. **We expect that all available monoclonal antibodies available in Switzerland are currently ineffective against the majority of the circulating SARS-CoV-2 viruses.**

In order to get a faster turnaround relative to sequencing, 346T specific PCR was implemented in Geneva. At the end of November, 65.6% of sequences contained the 346T mutation, with another 6.1% of results being indeterminate. As of week 50, over 80% of sequences in Geneva contained this mutation which effectively confers complete escape from Evusheld

neutralization (Table 2). Note that other mutations at position 346 as well as 444 also confer complete escape from neutralization by Evusheld, and the percent of sequences bearing the 346T mutation represent a minimum estimate for the number of sequences corresponding to virus that completely escape neutralization by Evusheld.

Date (week)	Samples	Number 346T	Number Indeterminate	% R346T	% Indeterminate
31.10.-06.11. (44)	80	42	7	52.5	13.3
07.11.-13.11. (45)	86	55	9	64.0	14.1
14.11.-20.11. (46)	94	66	6	70.2	8.5
21.11.-27.11 (47)	93	61	4	65.6	6.1
28.11.-4.12 (48)	94	74	1	78.7	1.3
5.12.-11.12 (49)	93	74	4	79.6	5.0
12.12.-18.12 (50)	93	81	1	87.1	1.1


*Table 2: Results of a PCR test specific for the R346T mutation run on samples taken at the Geneva university hospitals. The melting curve analysis of some samples had results outside of the normal range for unmutated 346, but that did not match the normal melting curve for a 346T mutation. These samples were considered "indeterminate", and may represent a mutation to another residue at position 346T. Also note that some mutations at position 346 still have a melting curve indistinguishable from an unmutated position 346.*

### **Epidemiology in Switzerland and number and origin of sequences produced through the program during the surveilled period**

Data in this report comes from 3 sources: 1) The publicly available data on COVID-19 as reported by the FOPH (<https://www.covid19.admin.ch>), including data that is declared to the FOPH by the different laboratories in Switzerland; 2) data originating from laboratories participating in the surveillance program; and 3) sequences submitted to GISAID, for which the corresponding infected person was in Switzerland (resident or recent travel history to Switzerland).

General caveat: the numbers and denominators are fluid and variable over time; and are subject to change depending notably on the different databases used, and variable declaration delays. All data generated by this program is also submitted to SPSP.

Data will be presented here by regions, using the same region definitions that are used for the influenza sentinel surveillance system in Switzerland. Data are presented according to residency post-code.



Region 1 includes the cantons of Geneva, Neuchatel, Vaud and Wallis

Region 2 includes the cantons of Bern, Fribourg and Jura

Region 3 includes the cantons of Aargau, Basel (Basel-Stadt and Basel-Land) and Solothurn

Region 4 includes the cantons of Luzern, Unterwalden (Obwalden and Nidwalden), Schwitz, Uri and Zug

Region 5 includes the cantons of Appenzell (Appenzell Ausserrhoden and Appenzell Innerrhoden), Glarus, Sankt Gallen, Schaffhausen, Thurgau and Zurich.

Region 6 includes the cantons of Graubünden and Ticino.

Divisions of the different regions, from <https://covariants.org/per-country>

### **Number of cases processed by the laboratories participating in the surveillance program**

During November (31 October to 27 November), the FOPH reported a total of 82'461 confirmed SARS-CoV-2 cases in Switzerland (according to data received by 23-12-2022), representing a substantial decrease from October (127'053 cases). Supplementary Table 1 provides an overview of the number and incidence of confirmed cases, the effective reproduction number  $R_e$ , the number and incidence of tests, test positivity, the number and proportion of sequenced samples, and the number and proportion of VOCs by canton, region and for Switzerland overall.

As of the writing of this report, the laboratories participating in this program reported 24'624 positive tests during the surveilled program, which represents 29.9% of the total number of cases reported in Switzerland (including both PCR and antigen-based tests), which very close to the 29.2% figure of October.

Detailed data regarding the total number of tests performed each week by the laboratories participating in the surveillance program (including negative and positive tests numbers, and the number of the positive tests that have been sequenced) are available in supplementary Table 3.

**Number of declared SARS-CoV-2 sequences produced through the surveillance program (presented by submission date, further declarations are still ongoing)**

A total of 2'109 SARS-CoV-2 sequences have been declared to have been submitted to GISAID during this period. There are 2'131 sequences available that were collected during this period on GISAID (and 2'143 submitted during this period) as of 22 December 2022, and the difference may be explained by reporting delays and differences between collection/submission dates. This represents around 2.6% of all cases detected in Switzerland during the surveilled period (see Supplementary Table 2 and 3 in the Appendix for details), an increase from October. This reflects a similar sequencing volume and declining case numbers. Of note, this number includes sequences from samples received from other laboratories at the request of the Cantonal physician teams.

Table 3 shows the number of sequences successfully submitted to GISAID through the surveillance program during the surveilled period by calendar week.


Week	Date	Number of sequences declared and successfully submitted to GISAID during the surveilled period, by all laboratories in the program
44	Oct. 31 to Nov. 6	486
45	Nov. 7 to Nov. 13	639
46	Nov. 14 to Nov.20	392
47	Nov. 21 to Nov. 27	592
<b>Total</b>		<b>2'109</b>

*Table 3: number of sequences submitted to GISAID through the surveillance program. Note these data are not by sampling date but rather by submission to GISAID date.*

The total number of SARS-CoV-2 sequences declared and submitted to GISAID by each laboratory during this period is available in Supplementary Table 3 in the appendix.

**Covering of sequencing in Switzerland and contribution of the national SARS-CoV-2 surveillance sequencing program**

As shown in Figure 1, the total number of SARS-CoV-2 sequences submitted per week generally remained stable during November (weeks 44-47), while the fraction sequenced was higher than in October. Since the beginning of this program, almost all of the sequences available, and on which the surveillance is conducted, come from the national surveillance program.



*Figure 1: Number of SARS-CoV-2 sequences available for Switzerland (total available Swiss sequences in GISAID in green, Swiss sequences submitted through the program in dotted orange) and fraction of the total number of positive cases declared to the FOPH that have been sequenced (blue curve).*



During the surveilled period, the absolute number of sequences generated remained fairly high, at hundreds of sequences per week. The total proportion of positive sequenced cases was approximately 2.6% during the month. These sequences include those from sites with hospitalized patients, and thus should be adequate for surveillance.

Figure 2 displays the number and fraction of SARS-CoV-2 cases sequenced for each Swiss region. Regions 4 and 6 had the lowest total number of sequences, while region 4 had the lowest fraction of cases sequenced.




Figure 2: Sequencing coverage among the different Swiss regions per week, presented by fraction of cases sequenced (left) and by number of sequences (right).

#### 4. Recently circulating variants in Switzerland as of August 2022

Determination of the proportion of total number of sequences over time falling into defined variant groups is done by Emma Hodcroft's team and displayed on the CoVariant website (<https://covariants.org/per-country>). Those results are based on the total number of sequences submitted to GISAID over the time period for Switzerland. Those data mainly, but not exclusively, come from the national genomic surveillance program since its beginning (see Figure 1).

Basal Omicron BA.5 was narrowly surpassed by its BQ.1 sublineage (Figures 3-5, Table 4) as the most common variant in Switzerland during November. Delta and BA.1 were not detected in Switzerland in November. An estimate of the total number of VOCs circulating in Switzerland, corrected by taking in account the fraction of sequencing in Switzerland is available through the covSPECTRUM program, developed at ETHZ, at <https://cov-spectrum.ethz.ch/explore/Switzerland>. Notably, 113 BA.2.75 sequences were found in Switzerland in the month of November, an increase from October. BA.2.75 sequences are expected to be over represented due to preferentially sequencing of S-gene dropout samples.

Region	BA.2*	BA.2.75*	BA.4*	BA.5*	BQ.1*	other	sequences	cases	% sequenced
All	20	113	26	632	940	400	2131	82461	2.58
1	3	14	10	152	372	73	624	15941	3.91
2	8	59	3	110	149	87	416	15850	2.625
3	2	5	1	138	148	106	400	15407	2.60
4	1	0	0	18	13	7	39	6120	0.64
5	6	31	9	161	174	102	483	22816	2.12
6	0	3	3	47	62	23	138	6327	2.18

Table 4: number of sequences corresponding to selected variants in each region of Switzerland from 31 October 2022 to 27 November 2022, according to data received by 23 December, 2022.






Figure 3: proportion of the total number of sequences (not cases), over time, that fall into defined variant groups, for Switzerland. Screenshot from CoVariants website. Dynamic navigation is available at <https://covariants.org/per-country>. Dark green indicates the B.1.617.2 (Delta) lineage or its sub-lineages. Dark Red indicates B.1.1.7 (Alpha). Purple/21K indicates Omicron BA.1, Light purple/21L indicates Omicron BA.2. Dark blue/22C indicates Omicron BA.2.12.1, while a faint blue/22B indicates Omicron BA.5 and a blueish-purple/22A indicates Omicron BA.4. Cyan/22D indicates BA.2.75. Bright purple (22E) indicates BQ.1, while 22F indicates the recombinant XBB lineage




Figure 4: (Left): Percentage of circulating VOCs and VOIs in Switzerland by week, up to the 47 first weeks of 2022 (total number of BA.2, BA.2.75, BA.4, BA.5, and BQ.1 sequences from Switzerland and successfully submitted to GISAID are shown here). Note the grey shaded area indicates a period of incomplete data. (Right): Estimated number of sequences BA.2, BA.2.75, BA.4, BA.5, and BQ.1 retrieved during the surveilled period.





Figure 5: Distribution of variants per region, for the end of November 2022 shown on a map. The size of the pie chart corresponds to the total number of sequences. Note the large share of the share of the BQ.1\* lineage, and the significant fraction of BA.2.75\* variants in some Swiss regions.

## 5. Assessment of the competition between the different variants in Switzerland

The competition between different SARS-CoV-2 variants can be modelled using multinomial logistic regression. The analysis by Dr. Althaus' group is based on sequences retrieved from CovSPECTRUM. The results correctly predicted that Omicron would be the dominant SARS-CoV-2 variant in Switzerland. These models also correctly predicted that the BA.1 sub-lineage would be replaced by the BA.2 sublineage (Figure 6). In May, the analysis predicted that BA.2 would be displaced by other variants (mainly BA.5), which was indeed correct. The current estimates suggest that basic BA.5 prevalence will continue to decline, with other lineages displacing it. The model predicts a rise of the recombinant XBB lineage, but not the BA.2.75 lineage.



*Figure 6: Observed and modeled proportion of SARS-CoV-2 variants over time in Switzerland. Since early 2021, multiple successive variants have been observed to displace all previous variants (often achieving a dominance of >99.9% of the circulation). Alpha was replaced by Delta, followed by Omicron BA.1, then Omicron BA.2, and now Omicron BA.5. Model fits are based on a multinomial logistic regression with splines.*

## 6. Surveillance of mutations associated with reduced mAb treatment efficacy

A number of mutations have been reported to result in significantly reduced *in vitro* effectiveness of the monoclonal antibodies used to treat patients in Switzerland. Notably, all Omicron lineages are completely resistant to neutralization by casirivimab/imdevimab. Sotrovimab retains substantial neutralizing activity against BA.1, but this neutralization is significantly decreased for BA.2/4/5 (and in particular, some of their sublineages). The escape is however not complete, and sotrovimab retains significant activity at higher titers – thus the frequencies of mutations (aside from lineage defining mutations) reported to escape neutralization by sotrovimab are still being followed (Table 5).

	337 (H, L, R, S, T)		340 (A, D, G, K, Q, V)		356 (T)		371 (F, L)	
Dates	Global	CH/EU	Global	CH/EU	Global	CH/EU	Global	CH/EU
31.10 to 27.11	0.01	0.01/0.01	0.03	0.09/0.05	4.37	4.96/4.71	95.19	99.25/93.83

	377 (K)		449 (N)		476 (S)		494 (P)	
Dates	Global	CH/EU	Global	CH/EU	Global	CH/EU	Global	CH/EU
31.10 to 27.11	0.00	0.00/0.00	0.0	0.00/0.00	0.08	0.00/0.06	0.25	0.66/0.25

Table 5: Percent (to the nearest 0.01%) of Global and Swiss sequences bearing Sotrovimab escape mutations (>5x neutralization drop or >0.8 escape fraction) for November 2022(31 October to 27 November)

Notably, additional escape mutations against Sotrovimab are currently rare in Switzerland and world-wide. However, the vast majority of circulating variants contain the 371 escape mutations, which are associated with escape >10x) from Sotrovimab.

Evusheld®, which consists of a cocktail of 2 mabs, tixagevimab and cilgavimab, is also used in Switzerland. Tixagevimab is not effective against BA.2/4/5, although the BA.2.75 sublineage is neutralized by tixagevimab, likely due to a reversion mutation. Cilgavimab is able to neutralize BA.2/4/5, although its neutralization is reduced against BA.4/5, and eliminated in the BA.4.6/4.7/5.9 sublineages.

Mutations resulting in partial escape from cilgavimab are shown in table 6. As seen in tables 5&6, in contrast to the situation with Sotrovimab, known mutations enabling escape from Evusheld® are increasing rapidly in Switzerland and world-wide. During November, variants expected to be completely resistant to Evusheld® (cilgavimab and tixagevimab) made up over 79% of the sequences, and preliminary data for December shows a proportion of nearly 90%. Many of these escape mutations are also associated with escape from humoral immunity, thus it is not expected to be seen only in response to mAb treatment.

	346 (I, N, S, T)		444 (E,G,L,M,N,Q,R,S,T)		445 (A)		446 (S)	
Dates	Global	CH/EU	Global	CH/EU	Global	CH/EU	Global	CH/EU
31.10 to 27.11	50.08	66.12/56.32	36.19	47.36/39.04	1.48	1.60/0.84	11.48	10.79/9.67

	447 (R, V)		448 (Y)		450 (D, K)		452 (R)	
Dates	Global	CH/EU	Global	CH/EU	Global	CH/EU	Global	CH/EU
31.10 to 27.11	0.00	0.00/0.00	0.00	0.00/0.00	2.60	1.93/3.03	83.95	87.32/85.86

	484 (A)		494 (R)		655 (Y)	
Dates	Global	CH/EU	Global	CH/EU	Global	CH/EU
31.10 to 27.11	95.46	98.92/94.64	0.00	0.00/0.00	98.78	99.67/98.65

Table 6: Percent of Global and Swiss sequences bearing cilgavimab escape mutations (>5x neutralization drop or >0.8 escape fraction) for November 2022(31 October to 27 November).

	371 (F)		417 (N)		475 (D)		476 (D)	
Dates	Global	CH/EU	Global	CH/EU	Global	CH/EU	Global	CH/EU
31.10 to 27.11	95.17	98.92/93.81	92.79	97.46/92.73	0.00	0.00/0.00	0.00	0.00/0.00

	484 (A, D, K)		486 (P, S, V)		493 (R)		498 (R)	
Dates	Global	CH/EU	Global	CH/EU	Global	CH/EU	Global	CH/EU
31.10 to 27.11	95.46	98.92/94.64	89.84	92.88/88.74	0.06	0.09/0.09	95.77	99.76/94.07

	990 (A)		1009 (I)	
Dates	Global	CH/EU	Global	CH/EU
31.10 to 27.11	0.00	0.00/0.00	0.05	0.05/0.05

Table 7: Percent of Global and Swiss sequences bearing Tixagevimab escape mutations (>5x neutralization drop or >0.8 escape fraction) for November 2022(31 October to 27 November). Note: defining mutations of the Dominant BA.4/5 Omicron lineages and are not shown.

Paxlovid®, which inhibits the main viral protease encoded by the viral gene nsp5 is also used in Switzerland. In the absence of any treatment with Paxlovid®, escape mutations are not expected to produce any benefit, and the mutations are not linked to general antigenic shift like the escape mutations to the therapeutic mAbs (ie: sotrovimab and Evusheld®). This likely explains the scarcity of escape mutations against Paxlovid®. Notably, while dozens of mutations at sites known to be important for escape from Paxlovid have been reported worldwide (table 1), only 2 mutations at those sites that are actually known to cause escape have been sequenced worldwide during November, (both from Europe (table 8), resulting in a miniscule percentage of total sequences. Mutations resulting in partial escape from Paxlovid® are shown in table 8.

	54 C		138 S		144 A/F/G/M/T/Y		165 T	
Dates	Global	CH/EU	Global	CH/EU	Global	CH/EU	Global	CH/EU
31.10 to 27.11	0	0	0	0	0	0	0	0

	166 Q/V		167 F		172 Q/F		192 R/S/T/V	
Dates	Global	CH/EU	Global	CH/EU	Global	CH/EU	Global	CH/EU
31.10 to 27.11	1 (V)	0/1 (V)	0	0	0	0	0	0

	194 S		305 L	
Dates	Global	CH/EU	Global	CH/EU
31.10 to 27.11	0	0	1	0/1

Table 8: Global and Swiss counts of sequences bearing escape mutations from Paxlovid®: Sequenced escape mutations remained extremely rare worldwide during the month of November (31 October to 27 November).

## 7. Wastewater surveillance program

Sequences are obtained from six wastewater treatment plants (WWTPs) that are tested on a daily basis. At the start of the month of November, BA.5 was still the major variant, making up 53% to 87% of the sequenced SARS-CoV-2 RNA in all locations (Figures 7, 8). During the month of November, B.Q.1.1, continued rising, approaching or surpassing 50% of the sequenced SARS-CoV-2 RNA, depending on the location. During the same month, the BA.2.75 variant was consistently detected in all surveyed treatment plants. At the end of November, it accounted for up to 8% of the sequenced SARS-CoV-2 RNA, depending on the location. The XBB variant has been consistently observed in multiple treatment plant since the beginning of November, but does not show sustained growth at the moment.




Figure 7:

Overview of the relative abundances of variants of SARS-CoV-2 at the end of November 2022, estimated from wastewater samples collected daily during the month in WWTPs located in 6 different Swiss locations. Pie chart areas are proportional to population connected to the wastewater treatment plants. Pink shaded areas represent catchment areas (boundaries from 2017). Population connected to Vernier (GE) wastewater treatment plant also includes ~44,000 inhabitants from neighbouring French communities.




Figure 8: Relative abundances of variants of SARS- CoV-2 estimated from wastewater samples collected daily until November 30, 2022 in WWTPs located in 6 different Swiss cantons. An online dynamic navigation is available at <https://bsse.ethz.ch/cbq/research/computational-virology/sarscov2-variants-wastewater-surveillance.html>.



**Acknowledgements:**

<https://bsse.ethz.ch/cevo/research/sars-cov-2/swiss-sars-cov-2-sequencing-consortium.html>

Meriem Bekliz, Kenneth Adea, Pauline Vetter, Christiane S Eberhardt, Krisztina Hosszu-Fellous, Diem-Lan Vu, Olha Puhach, Manel Essaidi-Laziosi, Sophie Waldvogel-Abramowski, Caroline Stephan, Arnaud G. L'Huillier, Claire-Anne Siegrist, Arnaud M Didierlaurent, Laurent Kaiser, Benjamin Meyer, Isabella Eckerle for the characterization neutralization of Omicron by patient sera.

Erik Boehm, Marc Friedli, Pauline Vetter, Samuel Cordey, Richard Neher, Christian Althaus, Emma Hodcroft, Tanja Stadler, Philippe Lemercier, Ioannis Xenarios, Lorenzo Cerutti, Louis Du Plessis, Erik Studer, Oluwaseun Oyewole, Laurent Kaiser, for the Swiss national SARS-CoV-2 genomic and variants surveillance program coordination committee.

We would also like to thank the CoVICIS project (<https://covicis.eu/>), which is funded by an EU grant, for supplementary funding for genomic sequencing in Geneva.

**Appendix:****SARS-CoV-2 epidemiology in Switzerland:**

We used publicly available data on COVID-19 as reported by FOPH (<https://www.covid19.admin.ch>) and sequence data submitted to GISAID to provide a summary of the SARS-CoV-2 epidemiology in Switzerland.



sup\_table\_overview  
\_Nov.xlsx

*Supplementary Table 1: Epidemiological data for Switzerland, its regions and cantons for November: population, number and incidence of confirmed cases, effective reproduction number  $R_e$ , number and incidence of tests, test positivity, number and proportion of sequenced samples, and number and proportion of VOCs.  $R_e$  by region is represented as the median and range of the daily  $R_e$  values for all cantons within a region.*

week	Date	Total PCR tests	Positive tests	Sequenced	% positives	% positives sequenced
44	Oct. 31 to Nov. 6	18'535	6'321	486	34.10%	7.69%
45	Nov. 7 to Nov. 13	18'856	6'038	639	32.02%	10.58%
46	Nov. 14 to Nov.20	19'639	6'312	392	32.14%	6.21%
47	Nov. 21 to Nov. 27	19'473	5'953	592	30.57%	9.94%
	<b>Total</b>	76'503	2'624	2'109	32.19%	8.56%

*Supplementary Table 2: Total number of tests performed by the laboratories participating in the surveillance program from 31 October to 27 November 2022.*

Week	Date	Basic Surveillance				Augmented Surveillance						Sentinella Laboratories		
		EOC	Labor Team W	Risch	SRO	USB	IFIK	Diana labs	CHUV	UZH	ICH-VS*	HUG	ETH/ Viollier	All
44	Oct. 31 to Nov. 6	18	19	59	7	0	74	44	38	100	39	89	73	486
45	Nov. 7 to Nov. 13	19	18	56	14	77			39	80		176	86	639
46	Nov. 14 to Nov.20	15	25	54	12	38	69	42	37	81	39	165	91	392
47	Nov. 21 to Nov. 27	21	21	55	10	39			55	40			75	592
	<b>Total</b>	73	83	224	43	154	143	86	169	301	78	430	325	2'109

*Supplementary Table 3: number of sequences submitted to GISAID by each laboratory during the surveilled period (31 October to 27 November 2022). \*including sequencing sent to high-throughput platforms.*

## Contact list as of 22.07.22 :

Coordination committee mailing list	
Name	e-mail address
Laurent Kaiser	<a href="mailto:Laurent.Kaiser@hcuge.ch">Laurent.Kaiser@hcuge.ch</a>
Samuel Cordey	<a href="mailto:Samuel.Cordey@hcuge.ch">Samuel.Cordey@hcuge.ch</a>
Marc Friedli	<a href="mailto:marc.friedli@epfl.ch">marc.friedli@epfl.ch</a>
Richard Neher	<a href="mailto:richard.neher@unibas.ch">richard.neher@unibas.ch</a>
Tanja Stadler	<a href="mailto:tanja.stadler@bsse.ethz.ch">tanja.stadler@bsse.ethz.ch</a>
Louis Du Plessis	<a href="mailto:louis.duplessis@bsse.ethz.ch">louis.duplessis@bsse.ethz.ch</a>
Emma Hodcroft	<a href="mailto:emma.hodcroft@ispm.unibe.ch">emma.hodcroft@ispm.unibe.ch</a>
Christian Althaus	<a href="mailto:christian.althaus@ispm.unibe.ch">christian.althaus@ispm.unibe.ch</a>
Ioannis Xenarios	<a href="mailto:ioannis.xenarios@unil.ch">ioannis.xenarios@unil.ch</a>
Philippe Le Mercier	<a href="mailto:Philippe.Lemercier@sib.swiss">Philippe.Lemercier@sib.swiss</a>
Pauline Vetter	<a href="mailto:Pauline.Vetter@hcuge.ch">Pauline.Vetter@hcuge.ch</a>
Erik Boehm	<a href="mailto:Erik.Boehm@hcuge.ch">Erik.Boehm@hcuge.ch</a>
Lorenzo Cerutti	<a href="mailto:lorenzo.cerutti@health2030.ch">lorenzo.cerutti@health2030.ch</a>
Erik Studer	<a href="mailto:Erik.Studer@bag.admin.ch">Erik.Studer@bag.admin.ch</a>
Oluwaseun Oyewole	<a href="mailto:Oluwaseun.Oyewole@bag.admin.ch">Oluwaseun.Oyewole@bag.admin.ch</a>

Laboratories mailing list		
Laboratory	Name	e-mail address
HUG	Laurent Kaiser	<a href="mailto:Laurent.Kaiser@hcuge.ch">Laurent.Kaiser@hcuge.ch</a>
HUG	Samuel Cordey	<a href="mailto:Samuel.Cordey@hcuge.ch">Samuel.Cordey@hcuge.ch</a>
HUG	Pauline Vetter	<a href="mailto:Pauline.Vetter@hcuge.ch">Pauline.Vetter@hcuge.ch</a>
HUG	Erik Boehm	<a href="mailto:Erik.Boehm@hcuge.ch">Erik.Boehm@hcuge.ch</a>
CHUV	Gilbert Greub	<a href="mailto:Gilbert.Greub@chuv.ch">Gilbert.Greub@chuv.ch</a>
CHUV	Claire Bertelli	<a href="mailto:Claire.Bertelli@chuv.ch">Claire.Bertelli@chuv.ch</a>
Universtätsspital Basel	Pascal Schlaepfer	<a href="mailto:Pascal.Schlaepfer@usb.ch">Pascal.Schlaepfer@usb.ch</a>
Universtätsspital Basel	Karoline Leuzinger	<a href="mailto:Karoline.Leuzinger@usb.ch">Karoline.Leuzinger@usb.ch</a>
Universtätsspital Basel	Hans Hirsch	<a href="mailto:Hans.Hirsch@usb.ch">Hans.Hirsch@usb.ch</a>
IFIK UNIBE	Alban Ramette	<a href="mailto:alban.ramette@ifik.unibe.ch">alban.ramette@ifik.unibe.ch</a>
UZH	Alexandra Trkola	<a href="mailto:trkola.alexandra@virology.uzh.ch">trkola.alexandra@virology.uzh.ch</a>
UZH	Michael Huber	<a href="mailto:huber.michael@virology.uzh.ch">huber.michael@virology.uzh.ch</a>
EOC Bellinzona	Gladys Martinetti Luchini	<a href="mailto:Gladys.MartinettiLucchini@eoc.ch">Gladys.MartinettiLucchini@eoc.ch</a>
Stadler group/Viollier laboratories	Louis du Plessis	<a href="mailto:louis.duplessis@bsse.ethz.ch">louis.duplessis@bsse.ethz.ch</a>
Stadler group/Viollier laboratories	Tanja Stadler	<a href="mailto:tanja.stadler@bsse.ethz.ch">tanja.stadler@bsse.ethz.ch</a>
Viollier laboratories	Christiane Beckmann	<a href="mailto:christiane.beckmann@viollier.ch">christiane.beckmann@viollier.ch</a>
Viollier laboratories	Henriette Kurth	<a href="mailto:Henriette.Kurth@viollier.ch">Henriette.Kurth@viollier.ch</a>
Hopitaux du Valais – Institut Central	Alexis Dumoulin	<a href="mailto:Alexis.Dumoulin@hopitalvs.ch">Alexis.Dumoulin@hopitalvs.ch</a>
Dianalabs	Nadia Liassine	<a href="mailto:Nadia.liassine@dianalabs.ch">Nadia.liassine@dianalabs.ch</a>
Dianalabs	Katia Jaton	<a href="mailto:Katia.jaton@dianalabs.ch">Katia.jaton@dianalabs.ch</a>
Dianalabs	Géraldine Jost	<a href="mailto:Geraldine.jost@dianalabs.ch">Geraldine.jost@dianalabs.ch</a>
Dianalabs (Genesupport)	Tanguy Araud	<a href="mailto:Tanguy.araud@genesupport.ch">Tanguy.araud@genesupport.ch</a>
Laboratoire Bioanalytica	Michael Naegele	<a href="mailto:michael.naegele@bioanalytica.ch">michael.naegele@bioanalytica.ch</a>
Laboratoire Bioanalytica	Livia Berlinger	<a href="mailto:livia.berlinger@bioanalytica.ch">livia.berlinger@bioanalytica.ch</a>
Labor Team W ag	Andreas Lindauer	<a href="mailto:andreas.lindauer@team-w.ch">andreas.lindauer@team-w.ch</a>
Spital Region Oberaargau	Alexander Imhof	<a href="mailto:a.imhof@sro.ch">a.imhof@sro.ch</a>
Laboratory Risch	Nadia Wohlwend	<a href="mailto:nadia.wohlwend@risch.ch">nadia.wohlwend@risch.ch</a>

BAG mailing list:	
Name	e-mail address
Erik Studer	<a href="mailto:Erik.Studer@bag.admin.ch">Erik.Studer@bag.admin.ch</a>
Oluwaseun Oyewole	<a href="mailto:Oluwaseun.Oyewole@bag.admin.ch">Oluwaseun.Oyewole@bag.admin.ch</a>
Katrin Schneider	<a href="mailto:katrin.schneider@bag.admin.ch">katrin.schneider@bag.admin.ch</a>
Anna Fesser	<a href="mailto:Anna.Fesser@bag.admin.ch">Anna.Fesser@bag.admin.ch</a>
Ursina Roder	<a href="mailto:ursina.roder@bag.admin.ch">ursina.roder@bag.admin.ch</a>
Lionel Rohner	<a href="mailto:lionel.rohner@bag.admin.ch">lionel.rohner@bag.admin.ch</a>
Michael Bel	<a href="mailto:Michael.Bel@bag.admin.ch">Michael.Bel@bag.admin.ch</a>
Urs Mayr	<a href="mailto:urs.mayr@bag.admin.ch">urs.mayr@bag.admin.ch</a>
Damir Perisa	<a href="mailto:Damir.Perisa@bag.admin.ch">Damir.Perisa@bag.admin.ch</a>
Nadia Corazza	<a href="mailto:nadia.corazza@bag.admin.ch">nadia.corazza@bag.admin.ch</a>
Martine Bourqui	<a href="mailto:Martine.Bourqui@bag.admin.ch">Martine.Bourqui@bag.admin.ch</a>
Natalia Krempaska	<a href="mailto:natalia.krempaska@bag.admin.ch">natalia.krempaska@bag.admin.ch</a>
Selina Schwegler	<a href="mailto:Selina.schwegler@bag.admin.ch">Selina.schwegler@bag.admin.ch</a>
Mirjam Mäusezahl	<a href="mailto:Mirjam.Mäusezahl@bag.admin.ch">Mirjam.Mäusezahl@bag.admin.ch</a>
Tobias Schuster	<a href="mailto:tobias.schuster@bag.admin.ch">tobias.schuster@bag.admin.ch</a>

Sequencing centers:		
Center	Name	e-mail address
Health 2030 Genome Center	Keith Harshman	<a href="mailto:keith.harshman@health2030.ch">keith.harshman@health2030.ch</a>
Health 2030 Genome Center	Ioannis Xenarios	<a href="mailto:ioannis.xenarios@health2030.ch">ioannis.xenarios@health2030.ch</a>
Genomics Facility Basel-ETH Zurich	Christian Beisel	<a href="mailto:christian.beisel@bsse.ethz.ch">christian.beisel@bsse.ethz.ch</a>
The Functional Genomics Center Zurich (FGCZ)- ETHZ and UZH	Ralph Schlapbach	<a href="mailto:ralph.schlapbach@fgcz.ethz.ch">ralph.schlapbach@fgcz.ethz.ch</a>

Wastewater surveillance program mailing list:	
Name	e-mail address
Niko Beerenwinkel	<a href="mailto:niko.beerenwinkel@bsse.ethz.ch">niko.beerenwinkel@bsse.ethz.ch</a>
David Dreifuss	<a href="mailto:david.dreifuss@bsse.ethz.ch">david.dreifuss@bsse.ethz.ch</a>

Immunological characterization program of the variant mailing list:	
Name	e-mail address
Priscilla Turelli	<a href="mailto:priscilla.turelli@epfl.ch">priscilla.turelli@epfl.ch</a>
Didier Trono	<a href="mailto:didier.trono@epfl.ch">didier.trono@epfl.ch</a>